

First evidence of neonicotinoid residues in a long-distance migratory raptor, the European honey buzzard (*Pernis apivorus*)

Article (Accepted Version)

Byholm, Patrik, Mäkeläinen, Sanna, Santangeli, Andrea and Goulson, Dave (2018) First evidence of neonicotinoid residues in a long-distance migratory raptor, the European honey buzzard (*Pernis apivorus*). *Science of The Total Environment*, 639. pp. 929-933. ISSN 0048-9697

This version is available from Sussex Research Online: <http://sro.sussex.ac.uk/id/eprint/76691/>

This document is made available in accordance with publisher policies and may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the URL above for details on accessing the published version.

Copyright and reuse:

Sussex Research Online is a digital repository of the research output of the University.

Copyright and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable, the material made available in SRO has been checked for eligibility before being made available.

Copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

1 **First evidence of neonicotinoid residues in a long-**
2 **distance migratory raptor, the European honey**
3 **buzzard (*Pernis apivorus*)**

4 Patrik Byholm^{a*}, Sanna Mäkeläinen^b, Andrea Santangeli^b & Dave Goulson^c

5

6 ^a Novia University of Applied Sciences, Raseborgsvägen 9, FI-10600 Ekenäs, Finland

7 ^b Finnish Museum of Natural History Luomus, P.O. Box 17 (P. Rautatiekatu 13), FI-00014

8 University of Helsinki, Finland

9 ^c School of Life Sciences, University of Sussex, BN1 9RH Brighton, United Kingdom

10

11 ***Corresponding author:** patrik.byholm@novia.fi

12

13 **Abstract**

14 The evidence of negative impacts of agricultural pesticides on non-target organisms is constantly
15 growing. One of the most widely used group of pesticides are neonicotinoids, used in treatments of
16 various plants, e.g. oilseed crops, corn and apples, to prevent crop damage by agricultural insect
17 pests. Treatment effects have been found to spill over to non-target insects, such as bees, and more
18 recently also to other animal groups, among them passerine birds. Very little is known, however, on
19 the presence of neonicotinoids in other wild species at higher trophic levels. We present results on
20 the presence of neonicotinoid residues in blood samples of a long-distant migratory food-specialist
21 raptor, the European honey buzzard. Further, we investigate the spatial relationship between
22 neonicotinoid residue prevalence in honey buzzards with that of crop fields where neonicotinoids
23 are typically used. A majority of all blood samples contained neonicotinoids, thiacloprid accounting
24 for most of the prevalence. While neonicotinoid residues were detected in both adults and nestlings,
25 the methodological limit of quantification was exceeded only in nestlings. Neonicotinoids were
26 present in all sampled nests. Neonicotinoid presence in honey buzzard nestlings' blood matched
27 spatially with the presence of oilseed plant fields. These are the first observations of neonicotinoids
28 in a diurnal raptor. For better understanding the potential negative sub-lethal of neonicotinoids in
29 wild vertebrates, new (experimental) studies are needed.

30

31 **Keywords**

32 agricultural pesticides; European honey buzzard *Pernis apivorus*; neonicotinoids; oilseed rape;
33 migratory raptor; turnip rape

34

1. Introduction

There is accumulating evidence on the negative impacts of agricultural pesticides aimed at particular groups of herbivorous insect pests on non-target organisms (Geiger et al., 2010; Goulson, 2013). For example, declines of grassland bird populations in the United States have been found to be more strongly linked to large-scale use of pesticides rather than to land-use change and intensification in farmland (Mineau and Whiteside, 2013). Similarly, global declines of pollinator species have been related with the extensive use of neonicotinoids, a widely used class of agricultural pesticides globally (Tsvetkov et al., 2017; Woodcock et al., 2017).

Neonicotinoids are used mainly in seed coating of a wide variety of cultivated plants from corn (*Zea mays*) to oilseed rape (*Brassica napus oleifera*) (Jeschke et al., 2011; Simon-Delso et al., 2015). Targeted at sucking and boring insect pests, such as aphids and wireworm larvae, they effectively bind to the neural receptors of insects eventually causing paralysis and death (Tomizawa and Casida, 2003). However, much of the active ingredient does not end up in the crop but instead contaminates soil (Goulson, 2013; Jones et al., 2014), water and non-target foliage, including wild flowers growing in farmland (Botías et al., 2015; David et al., 2016). Neonicotinoids have been found to represent a major threat to bees through increased mortality and decreased colony establishment (e.g. Tsvetkov et al., 2017; Woodcock et al., 2017). Reports on adverse effects in other invertebrate groups are also accumulating (Pisa et al., 2017). It was long assumed that neonicotinoids have negligible impacts on vertebrate species due to their lower toxicity, yet recent studies have reported adverse effects of neonicotinoids on both terrestrial and aquatic vertebrates, even when found in concentrations well below the level causing acute poisoning or lethality (Crosby et al., 2015; Gibbons et al., 2015). To this end, subtle consequences such as impaired migratory ability, decreased body condition and breeding success in granivorous birds following the ingestion of small amounts of neonicotinoid coated grains have been shown (Lopez-Antia et al., 2015; Millot et al., 2017). Moreover, the toxicity level of neonicotinoids may vary greatly among

60 species, and may differentially impact on all or some of the key life-stages (e.g. breeding, survival,
61 migration) of a species (e.g. Gibbons et al., 2015; Eng et al., 2017). In the case of small bird
62 species, even ingestion of a single treated grain can cause acute intoxication and cause adverse
63 effects (Mineau and Palmer, 2013). Moreover, indirect effects of neonicotinoids causing a lack of
64 insects and other invertebrate food will lead to food deprivation in many species that depend on
65 invertebrates for food (cf. Hallman et al. 2014).

66 In the terrestrial realm, most scientific and political attention on the adverse effects of
67 neonicotinoids has largely focused on species groups that are in direct contact with neonicotinoid-
68 treated crops, such as invertebrates and granivorous birds (EU, 2013; Gibbons et al., 2015; Eng et
69 al., 2017). However, as neonicotinoids are known to spillover into the environment and the food-
70 chain in which they are introduced, there is a risk they may be transported towards the higher levels
71 of the food chain. However, to date investigations of neonicotinoids spilling over to the top of the
72 food-chain, e.g. raptors, is extremely rare (but see Taliansky-Chamudis et al., 2017). There is
73 therefore need for further scientific investigations on the exposure to neonicotinoids of species at
74 different trophic levels of the food-chain beyond those in more direct contact with these pesticides.

75 Here we report on the prevalence of neonicotinoid residues in European honey buzzards
76 (*Pernis apivorus*; from here onward honey buzzard) breeding in Finland during the Boreal summer.
77 This is a relevant study species because it mainly feeds on the larvae of social Apoidea, especially
78 wasps (*Vespidae*) but also pollinating bumble-bees (*Bombus* sp.), which in turn feed on mass
79 flowering oilseed crops that are often treated with neonicotinoids (Ketola et al., 2015). Being a
80 long-distance migrant, the honey buzzard could also be exposed to neonicotinoids across its passage
81 range as well as on its wintering range in Sub-Saharan Africa. Residues of neonicotinoids have been
82 in fact found in honey collected in countries along the flyway of this raptor species (cf. Mitchell et
83 al., 2017). The objective of this study is to provide the first evaluation of prevalence of
84 neonicotinoids in a wild population of a food-specialist raptor species. To account for local

variation in the occurrence of oilseed plant fields (potential neonicotinoid source) between sample sites, the spatial match between neonicotinoid prevalence in sampled birds with that of oilseed rape and turnip rape (*Brassica rapus oleifera*) fields within the hunting range of breeding birds was also investigated.

2. Methods

The honey buzzard is a forest-dwelling migratory raptor breeding in the Palearctic and wintering in Sub-Saharan Africa (Cramp, 1980). In the northern hemisphere, during the breeding season it feeds almost entirely on larvae of wasps and bumblebees, while frogs and small birds constitute alternative food (Itämiä and Mikkola, 1972; Gamauf, 1999). Similarly, on the African wintering sites, insects constitute an important food source for honey buzzards (Cramp, 1980).

As a part of a long-term study on forest raptors (Byholm and Nikula, 2007; Vansteelant et al., 2017), honey buzzard nests were sought for opportunistically. From a subset of found nests situated in Norway spruces (*Picea abies*) at 6-12 m height, ten honey buzzards from five families (Table 1) were sampled for blood in Western Finland (latitude 61°14'–63°12' N, longitude 21°16'–23°31' E) during the nestling phase in July-August 2013 when the nestlings were approximately 25-35 days old. To minimize disturbance, nest visits involving handling of nestlings only never lasted longer than 45 minutes, but when adults were caught (see below) visits lasted for up to two hours. Approximately 0.1 ml blood was collected from the brachial vein of one breeding pair and eight nestlings (two being offspring of the same pair) using a needle and a small glass capillary. The blood was placed in an Eppendorf-tube, and moved to a freezer bag containing dry ice for transportation to a super freezer (-80° C). Four months post-sampling the blood samples were sent to the University of Sussex for analyses of the residues of the neonicotinoids acetamidprid (ACE), imidacloprid (IMC) and thiacloprid (THC). Analysis of neonicotinoid residues in samples was

109 performed using ultra high-performance liquid chromatography tandem mass spectrometry
110 (UHPLC-MS/MS) (Waters Acquity UHPLC system, Waters, Manchester, UK). Procedures are
111 described in full in David et al. (2015). Analyte concentrations in blank workup samples were all
112 below the method limit of detection (LOD).

113 As a part of ongoing work on honey buzzard movement ecology (Vansteelant et al., 2017; P.
114 Byholm et al., unpublished), the parental birds (n= 4) at two nests (nest #1 and #5, cf. Table 1) were
115 caught using a dho-gaza (Zuberogoitia et al., 2008) and equipped with solar-powered Argos-GPS
116 platform terminal transmitters (PTTs) (Microwave Telemetry Inc.) or UvABiTS-GPS-trackers
117 (Bouten et al., 2013) using body-loop harnesses made of Teflon ribbon (Kenward, 2004). Tags'
118 weight (22-25 g) corresponded to ca. 3% of the birds' body mass at deployment (833 ± 161 ;
119 $\text{avg} \pm \text{SD}$). The amount of delivered GPS-fixes delivered varied depending on tracker model and
120 programming. All work requiring special permits (visiting nests, handling of honey buzzards,
121 collection and storage of blood samples) was performed under special licenses issued by the
122 relevant Finnish authorities to PB (ESAVI/1592/04.10.03/2011, EPOELY/135/07.01.2013,
123 PIRELY/49/07.01/2013, VARELY/73/07.01/2013, VARELY/215/2015).

124 In order to investigate the spatial relationship between neonicotinoid prevalence and landscape
125 composition, we collated spatial data on the location of field parcels (Agency for Rural Affairs,
126 2012; 2013) representing crop types typically treated with neonicotinoids in the study region in
127 Finland, i.e. spring turnip rape and oilseed rape (Ketola et al., 2015). Next, we extracted the fields
128 used in cultivation of oilseeds for the year 2013 (i.e. the year when blood sample data on the birds
129 were collected). The area of fields with the above crop types were then calculated within circular
130 buffers of radius 500 m, 1 km, 2 km and 5 km centered on each of the five honey buzzard nest site.
131 Disregarding night fixes (10:00 PM-5:00 AM) and fixes ≤ 80 m from the nest, the distances
132 between GPS-locations delivered by the GPS-trackers and the nests were calculated for the four
133 adult honey buzzards during whole breeding season (May 20th-August 29th). This material was then

134 used as a proxy for understanding over which distances from the nest the parents move while
135 hunting for their young. Finally we investigated whether (a) the amounts of neonicotinoids in honey
136 buzzard nestlings were correlated with the area of oilseed plant fields around honey buzzard nests,
137 and (b) whether presence of neonicotinoids in nestlings' blood was related to the presence of the oil
138 plant fields at different spatial scales. The latter was performed by Fisher's exact test. Here, when
139 the combined area of oil plant field parcels within a certain radius was below one hectare (nest #1 at
140 500 m), this was approximated to zero, based on its assumed negligible effects given the area
141 considered. Spatial data handling was done in ArcGIS 10.1 SP1 for Desktop (ESRI, 2012) and
142 statistical tests in R software for statistical computing, version 3.2.2 (R Core Team, 2015).

143

144 **3. Results**

145 Neonicotinoid residues were detected in the majority (8/10) of blood samples, adults and young
146 combined, at the methodological limit of detection (LOD; Table 1). Of these, 60% contained
147 neonicotinoids above the methodological limit of quantification (LOQ), with the amount of
148 compounds ranging from 8.9 to 31 pg/ml (average of $14.6 \pm \text{SD } 11.5$, Table 1). Among the
149 nestlings ($n = 8$), residues of imidacloprid or thiacloprid exceeding LOD and LOQ were found in
150 seven and six individuals, respectively. Among the two adults, imidacloprid exceeded the LOD in
151 one individual, whereas no neonicotinoid residues were found in the blood of adults above the
152 LOQ. Thiacloprid accounted for most of the quantified neonicotinoids prevalence, whereas
153 imidacloprid was less common and residues of acetamiprid were totally absent. Neonicotinoids
154 were detected in all of the five nests (Table 1).

155 Oilseed plant fields around the five different nest sites showed considerable variation at all
156 landscape scales (500 m-5 km). Some nests completely lacked oilseed plant fields ≤ 2 km from the
157 nest, but all nests had oilseed plant fields within a 5-km distance (Table 1). The overall proportion

158 of oilseed plant fields over the whole circular buffer area considered was generally small (max. 7%
159 of the total area in each case).

160 No correlation between the absolute area, or the percentage, of oil plant fields and the
161 quantified neonicotinoid levels in honey buzzard blood was found at any spatial scale (Pearson
162 correlation coefficients ranging from -0.11 to -0.27 and *p*-values from 0.46 to 0.78). However, when
163 neonicotinoid residues (of any compound) found in nestlings and the coverage of oil plant fields
164 around their nest were categorized into presence and absence, the presence/absence of
165 neonicotinoid residues matched with the presence/absence-pattern of oil plant fields at the scales of
166 2 and 5 km (Fisher's exact test, LOQ and LOD analyzed separately, all $p \geq 0.79$). No such spatial
167 match was found at the 1 km (LOQ: $p = 0.07$; LOD: $p = 0.02$) or 500-m scale (LOQ: $p = 0.007$,
168 LOD: $p = 0.007$). When comparing this result with the distance arrays describing how far from the
169 nest parental honey buzzards move when foraging, there is a fit with the spatial match observed for
170 neonicotinoid residue and oil plant field presence/absence. In other words, although adults differ in
171 respect to how far from the nest they forage, they typically forage at distances 2-5 km (or more)
172 than at 500m-1km from the nest (average median distance: 2.70 km; Figure 1).

173

174 **4. Discussion**

175 This is one of the first studies documenting the presence of neonicotinoid residues in a higher
176 trophic level consumer in a natural food web (but see Talianski-Chamudis et al., 2017). Of eight
177 blood sampled honey buzzard nestlings, residues of imidacloprid or thiacloprid exceeding LOD and
178 LOQ were found in seven and six individuals, respectively. LOD was also exceeded for
179 imidacloprid in the blood of one of the two sampled adults. No residues were found in the blood of
180 adults above the LOQ. Although there is limited knowledge about the persistence of neonicotinoids
181 in non-target animal species, the neonicotinoid *in vivo* metabolism observed in most animal species
182 is fast (Nishiwaki et al., 2004; Dick et al., 2005; Shao et al., 2013; Kapoor et al., 2014). Thus,

183 because neonicotinoids are water soluble (Goulson, 2013) and assuming the neonicotinoid
184 compounds in the honey buzzard blood circulation system in common with other species are
185 metabolized fast, the result that neonicotinoids were found in the majority of the blood samples of
186 nestlings implies that neonicotinoid exposure is of local origin rather than caught on the wintering
187 grounds or *en route* by adults during migration. If this inference is correct, it does not mean that an
188 individual testing negative for neonicotinoid residues could not have been in contact with the
189 pesticides during other parts of the annual cycle, but merely that testing for neonicotinoid residues
190 as sampled on the breeding grounds is an inadequate approach for identifying wintering ground
191 exposure. Thus, if aiming at understanding to what extent migratory birds are exposed to
192 neonicotinoids during the whole annual cycle, sampling must be organized regularly over wider
193 areas.

194 Although the sample size for this study is limited and absolute residue levels remain low, the
195 prevalence of neonicotinoid residues observed in this study is still higher than in previous studies
196 based on blood samples collected from other wild predators (Taliński-Chamudis et al., 2017).
197 Since honey buzzards prefer wasps and bumble-bees (*Bombus* sp.) as food, it seems most likely that
198 the reason for the neonicotinoid contamination is connected to this food source. Neonicotinoid
199 residues have commonly been found in the tissues of wild bees (Botías et al., 2017). Because
200 neonicotinoids in Finland in practice are used only to seed-coat and spray turnip rape and oilseed
201 rape (Ketola and Hakala, 2015), we can infer that the Apoidea species have been visiting these
202 fields for foraging and have come in direct contact with the neurotoxic pesticides. This conclusion
203 is further supported by the finding that the presence of neonicotinoid residues in honey buzzard
204 nestlings' blood matches strongly with the presence of oilseed plant fields within the home range at
205 distances of 2-5 km from the nest. These distances include the area of the home-range that is most
206 intensively used by foraging adult honey buzzards during the nestling provisioning phase (Fig. 1).
207 Consequently, provisioning parents likely deliver food that is contaminated by neonicotinoid

208 pesticides to their chicks, thereby explaining the presence of these substances found in the blood of
209 nestlings.

210 Species at the top of the food-chain, such as raptors, are in general highly sensitive to
211 environmental change, including the increasing prevalence of environmental pollutants, due to the
212 well-known process of bioaccumulation (Newton, 1976). Pollution, particularly through the use of
213 pesticides aimed at increasing crop yields, is pervasive across the cropland areas worldwide, with
214 seed-coating pesticides, such as neonicotinoids, being used across many countries in Europe, Africa
215 and elsewhere (www.fao.org/faostat). Most of these regions are very important for long-distance
216 Palearctic migrant birds breeding in Europe, such as the honey buzzard. Although the residue levels
217 reported here for honey buzzards are low, it should be noted that even a low dose may have
218 negative long-term effects (Goulson, 2013; Rondeau et al., 2014). Populations of long-distance
219 Palearctic migrants have been recently shown to be facing rapid declines, often driven by
220 anthropogenic pressures such as habitat loss and degradation, and overexploitation, outside of their
221 breeding grounds in Europe (Vickery et al., 2014; Laaksonen and Lehikoinen, 2013). Threats often
222 act in synergy in their adverse impacts on species (Brook et al., 2008). Exposure to pesticides, such
223 as neonicotinoids, may thus represent an additional pressure imposed on already declining
224 migratory species. Such added pressure may act on the non-breeding ground of the life of migratory
225 species (Eng et al., 2017), with potential consequences on survival, or on the breeding ground,
226 impacting on reproductive success (e.g. Lopez-Antia et al., 2015; Millot et al., 2017).

227 The findings presented here reinforce the scarce evidence so far available that neonicotinoid
228 substances can move up the top levels of the food-chain. However, neonicotinoid exposure of
229 predator species remains largely unquantified, and so are its possible consequences. Because the
230 rate of metabolism of neonicotinoids in birds is poorly understood, we cannot predict the dose of
231 neonicotinoids these honey buzzards were exposed to. Impacts of neonicotinoid exposure on birds
232 have been identified, ranging from direct mortality (Lopez-Antia et al., 2015; Millot et al., 2017) to

233 impaired migratory ability (Eng et al., 2017), compromised body condition and breeding success.
234 All such impacts are plausible to be present, and can lead to cascade effects, on species at the top of
235 the food-chain that become exposed to such pesticides. While our results are based on a small
236 sample size and cannot be generalized, they should sound as an early warning. We thus call the
237 scientific community to engage in collecting samples and assessing the exposure and its
238 consequences on species at the top levels of the food chain, both on the breeding and wintering
239 grounds. This endeavor is likely to be challenging methodologically, and would require long-term
240 studies to ultimately quantify the impacts of exposure on the population demography of long-lived
241 species. A first and relevant step, however, would be to quantify the extent of the potential problem.
242 That is, what species may be exposed to neonicotinoids. This study contributes to fill the above gap
243 in knowledge.

244 **Acknowledgements**

245 Our sincere thanks to Mikko Honkinieni, Ismo Nousiainen and Kari Palo for help with locating
246 active honey buzzard nests. Thanks are due to Dr David Arthur and Prof Elizabeth Hill for
247 assistance with the pesticide residue analysis. The comments from two anonymous referees greatly
248 improved the original version of the manuscript. This study has been funded by Finnish Cultural
249 Foundation's Southern Ostrobothnia Regional Fund (to SM), the Swedish Cultural Foundation in
250 Finland (to PB) and by the Academy of Finland (grant n. 307909; to AS). The sponsors had no role
251 in the study design, writing of the manuscript and submission process.

252 **Conflict of interest**

253 Authors state that there is no conflict of interest.

254

255 **References**

- 256 Agency for Rural Affairs, 2012. Land parcel register.
257 Agency for Rural Affairs, 2013. Crop type classification of field parcels.

258 Bouten, W., Baaij, E.W., Shamoun-Baranes, J., Camphuysen, K.C.J., 2013. A flexible GPS tracking
 259 system for studying bird behaviour at multiple scales. *J. Ornithol.* 154: 571–580.

260 Botías, C., David, A., Horwood, J., Abdul-Sada, A., Nicholls, E., Hill, E., Goulson, D., 2015.
 261 Neonicotinoid residues in wildflowers, a potential route of chronic exposure for bees. *Environ.*
 262 *Sci. Technol.* 49: 12731–12740.

263 Botías, C., David, A., Hill, E.M., Goulson, D., 2017. Quantifying exposure of wild bumblebees to
 264 mixtures of agrochemicals in agricultural and urban landscapes. *Environ. Pollut.* 222: 73–82

265 Brook, B.W., Sodhi, N.S., Bradshaw, C.J.A., 2008. Synergies among extinction drivers under
 266 global change. *Trends Ecol. Evol.* 23: 453–460.

267 Byholm, P., Nikula, A., 2007. Nesting failure in Finnish Northern goshawks *Accipiter gentilis*:
 268 Incidence and cause. *Ibis* 149: 597–604.

269 Cramp, S., 1980. Handbook of the Birds of Europe, the Middle East and North Africa: The Birds of
 270 the Western Palearctic. Vol. II. Oxford University Press, Oxford.

271 Crosby, E.B., Bailey, J.M., Oliveri, A.N., Levin, E.D., 2015. Neurobehavioral impairments caused
 272 by developmental imidacloprid exposure in zebrafish. *Neurotoxicol. Teratol.* 49: 81–90.

273 David, A., Botías, C., Abdul-Sada, A., Goulson, D., Hill, E.M. 2015. Sensitive determination of
 274 mixtures of neonicotinoid and fungicide residues in pollen and single bumblebees using a
 275 scaled down QuEChERS method for exposure assessment. *Analytical and Bioanalytical*
 276 *Chemistry* 407: 8151-8162.

277 David, A., Botías, C., Abdul-Sada, A., Nicholls, E., Rotheray, E.L., Hill, E.M., Goulson, D., 2016.
 278 Widespread contamination of wildflower and bee-collected pollen with complex mixtures of
 279 neonicotinoids and fungicides commonly applied to crops. *Environ. Int.* 88: 169–178.

280 Dick, R.A., Kanne, D.B., Casida, J.E., 2005. Identification of aldehyde oxidase as the neonicotinoid
 281 nitroreductase. *Chem. Res. Toxicol.* 18: 317–323.

282 Eng, M.L., Stutchbury, B.J.M., Morrissey, C.A., 2017. Imidacloprid and chlorpyrifos insecticides
 283 impair migratory ability in a seed-eating songbird. *Sci. Rep.* 7:15176. doi:10.1038/s41598-017-
 284 15446-x

285 ESRI, 2012. ArcGIS Desktop: Release 10.1. Redlands, CA: Environmental Systems Research
 286 Institute.

287 European Union, 2013. Commission Implementing Regulation (EU) No. 485/2013 amending
 288 Implementing Regulation (EU) No. 540/2011, as regards the conditions of approval of the
 289 active substances clothianidin, thiamethoxam and imidacloprid, and prohibiting the use and
 290 sale of seeds treated with plant protection products containing those active substances. *Official*
 291 *Journal of the European Union* 139: 12–26

292 Gamauf, A., 1999. Der Wespenbussard (*Pernis apivorus*) ein Nahrungsspezialist- Der Einfluß
 293 sozialer Hymenopteren auf Habitatnutzung und Home Range-Größe. *Egretta* 42: 57–85.

294 Geiger, F., Bengtsson, J., Berendse, F., Weisser, W.W., Emmerson, M., Morales, M.B., Ceryngier,
 295 P., Liira, J., Tscharntke, T., Winqvist, C., Eggers, S., Bommarco, R., Pärt, T., Bretagnolle, V.,
 296 Plantegenest, M., Clement, L.W., Dennis, C., Palmer, C., Oñate, J.J., Guerrero, I., Hawro, V.,
 297 Aavik, T., Thies, C., Flohre, A., Hänke, S., Fischer, C., Goedhart, P.W., Inchausti, P., 2010.
 298 Persistent negative effects of pesticides on biodiversity and biological control potential on
 299 European farmland. *Basic Appl. Ecol.* 11: 97–105.

300 Gibbons, D., Morrissey, C., Mineau P., 2015. A review of the direct and indirect effects of
 301 neonicotinoids and fipronil on vertebrate wildlife. *Environ. Sci. Pollut. Res.* 22: 103–118

302 Goulson, D., 2013. An overview of the environmental risks posed by neonicotinoid insecticides. *J.*
 303 *Appl. Ecol.* 50: 977–987.

304 Hallman, C., Foppen, R.P.B., van Turnhout, C.A.M., de Kroon, H., Jongelans, E., 2014. Declines in
 305 insectivorous birds are associated with high neonicotinoid concentrations. *Nature* 511: 341–
 306 343.

307 Itämies, J., Mikkola, H., 1972. The diet of Honey Buzzards *Pernis apivorus* in Finland. *Ornis Fenn.*
308 49: 7–10.

309 Jeschke, P., Nauen, R., Schindler, M., Elbert, A., 2011. Overview of the status and global strategy
310 for neonicotinoids. *J. Agric. Food Chem.* 59: 2897–2908.

311 Jones, A., Harrington, P., Turnbull, G., 2014. Neonicotinoid concentrations in arable soils after seed
312 treatment applications in preceding years. *Pest Manag. Sci.* 70: 1780–1784.

313 Kapoor, U., Srivastava, M.K., Trivedi, P., Garg, V., Srivastava, L.P., 2014. Disposition and acute
314 toxicity of imidacloprid in female rats after single exposure. *Food Chem. Toxicol.* 68: 190–95

315 Kenward, R., 2004. A manual for wildlife radio tagging. London, UK: Academic Press

316 Ketola, J., Hakala, K. 2015. Impact of use of neonicotinoid insecticides on honey bees in the
317 cultivation on spring oilseed crops in Finland. Interim report of the MTT Agrifood Research
318 Finland 157. 37 p.

319 Ketola J., Hakala K., Ruottinen L., Ojanen H., Rämö S., Jauhiainen L., Raiskio S., Kukkola M.,
320 Heinikainen S., Pelkonen S., 2015. The impact of the use of neonicotinoid insecticides on
321 honey bees in the cultivation of spring oilseed crops in Finland in 2013–2015. *Natural*
322 *resources and bioeconomy studies* 73/2015. Natural Resources Institute Finland (Luke). 70 p.

323 Laaksonen, T., Lehikoinen, A., 2013. Population trends in boreal birds: Continuing declines in
324 agricultural, northern, and long-distance migrant species. *Biol. Conserv.* 168: 99–107.

325 Lopez-Antia, A., Ortiz-Santaliestra, M.E., Mougeot, F., Mateo, R., 2015. Imidacloprid-treated seed
326 ingestion has lethal effect on adult partridges and reduces both breeding investment and
327 offspring immunity. *Environ. Res.* 136: 97–107. doi: 10.1016/j.envres.2014.10.023

328 Millot, F., Decors, A., Mastain, O., Quintaine, T., Berny, P., Vey, D., Lasseur, R., Bro, E., 2017.
329 Field evidence of bird poisonings by imidacloprid-treated seeds: a review of incidents reported
330 by the French SAGIR network from 1995 to 2014. *Environ. Sci. Pollut. Res.* 24: 5469–5485.

331 Mineau, P., Palmer, C., 2013. The Impact of the nation's most widely used insecticides on birds.
 332 American Bird Conservancy, March 2013.

333 Mineau, P., Whiteside, M., 2013. Pesticide acute toxicity is a better correlate of U.S. grassland bird
 334 declines than agricultural intensification. PLoS ONE 8(2).
 335 doi.org/10.1371/journal.pone.0057457

336 Mitchell, E.A.D., Mulhauser, B., Mullet, M., Mutabazi, A., Glauser, G., Aebi, A., 2017. A
 337 worldwide survey of neonicotinoids in honey. Science 358: 109–111.

338 Nishiwaki, H., Sato, K., Nakagawa, Y., Miyashita, M., Miyagawa, H., 2004. Metabolism of
 339 imidacloprid in houseflies. J. Pestic. Sci. 29: 110–116.

340 R Core Team, 2015. R: A language and environment for statistical computing. R Foundation for
 341 Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.

342 Newton, I., 1976. Population limitation in diurnal raptors. Can. Field Nat. 90: 274–300.

343 Pisa, L., Goulson, D., Yang, E.-C., Gibbons, D., Sánchez-Bayo, F., Mitchell, E., Aebi, A., van der
 344 Sluijs, J., MacQuarrie, C.J.K., Giorio, C., Long, E. Y., McField, M., Bijleveld van Lexmond,
 345 M., Bonmatin, J.-M., 2017. An update of the Worldwide Integrated Assessment (WIA) on
 346 systemic insecticides. Part 2: impacts on organisms and ecosystems. Environ. Sci. Pollut. Res.
 347 25: 1–49

348 Rondeau, G., Sánchez-Bayo, F., Tennekes, H.A., Decourtye, A., Ramírez-Romero, R., Desneux, N.,
 349 2014. Delayed and time-cumulative toxicity of imidacloprid in bees, ants and termites. Sci.
 350 Rep. 4: 5566.

351 Shao, X., Swenson, T.L., Casida, J.E., 2013. Cycloxaprid insecticide: nicotinic acetylcholine
 352 receptor binding site and metabolism. J. Agric. Food Chem. 61: 7883–7888

353 Taliansky-Chamudis, A., Gómez-Ramírez, P., León-Ortega, M., García-Fernández, A.J., 2017.
 354 Validation of a QuEChERS method for analysis of neonicotinoids in small volumes of blood

355 and assessment of exposure in Eurasian eagle owl (*Bubo bubo*) nestlings. Sci. Total Environ.
356 595: 93–100.

357 Tomisawa, M., Casida, J.E., 2003. Selective toxicity of neonicotinoids attributable to specificity of
358 insect and mammalian nicotinic receptors. Annu. Rev. Entomol. 48: 339–364.

359 Tsvetkov, N., Samson-Robert, O., Sood, K., Patel, H.S., Malena, D.A., Gajiwala, P.H.,
360 Maciukiewicz, P., Fournier, V., Zayed, A., 2017. Chronic exposure to neonicotinoids reduces
361 honey bee health near corn crops. Science 356: 1395–1397.

362 Vansteelant, W.M.G., Kekkonen, J., Byholm, P., 2017. Wind conditions and geography shape the
363 first outbound migration of juvenile honey buzzards and their distribution across sub-Saharan
364 Africa. Proc. R. Soc. B 284: 20170387.

365 Vickery, J. A., S. R. Ewing, K. W. Smith, D. J. Pain, F. Bairlein, J. Škorpilová, and R. D. Gregory.
366 2014. The decline of Afro-Palaearctic migrants and an assessment of potential causes. Ibis 156:
367 1–22.

368 Woodcock, B.A., Bullock, J.M., Shore, R.F., Heard, M.S., Pereira, M.G., Redhead, J., Ridding, L.,
369 Dean, H., Sleep, D., Henrys, P., Peyton, J., Hulmes, S., Hulmes, L., Sárospataki, M., Saure, C.,
370 Edwards, M., Genersch, E., Knäbe, S., Pywell, R.F., 2017. Country-specific effects of
371 neonicotinoid pesticides on honey bees and wild bees. Science 356: 1393–1395.
372 10.1126/science.aaa1190

373 Zuberogoitia, I., Martínez, J.E., Martínez, J.A., Zabala, J., Calvo, J.F., Azkona, A., Pagán, I., 2008.
374 The dho-gaza and mist net with Eurasian eagle-owl (*Bubo bubo*) lure: Effectiveness in
375 capturing thirteen species of European raptors, J. Raptor Res. 42: 48–51

376